



Role of pH level on the morphology and growth rate of myelin figures: publisher's note

MARZIEH ALLAH PANAHİ,^{1,4} ZAHRA TAHMASEBI,^{1,4} VAHID ABBASIAN,^{1,2,4} MOHAMMAD AMIRI,³ AND ALI-REZA MORADI^{1,2,5}

¹*Department of Physics, Institute for Advanced Studies in Basic Sciences (IASBS), Zanjan 45137-66731, Iran*

²*School of Nano Science, Institute for Research in Fundamental Sciences (IPM), Tehran 19395-5531, Iran*

³*Department of Physics, Bu-Ali Sina University (BASU), Hamedan 65175-4161, Iran*

⁴*These authors contributed equally to this work*

⁵*moradika@iasbs.ac.ir*

Abstract: This publisher's note contains corrections to *Biomed. Opt. Express* **11**, 5565 (2020).

© 2020 Optical Society of America under the terms of the [OSA Open Access Publishing Agreement](#)

This article [1] has been amended to include Fig. 2, which was cited in the paper but did not appear in the published paper. The paper was corrected online 13 October 2020.

References

1. M. A. Panahi, Z. Tahmasebi, V. Abbasian, M. Amiri, and A.-R. Moradi, "Role of pH level on the morphology and growth rate of myelin figures," *Biomed. Opt. Express* **11**(10), 5565–5574 (2020).